

Eligibility for Renal Denervation: Experience at 11 European Expert Centers

Alexandre Persu, Yu Jin, Marie Baelen, Eva Vink, Willemien L. Verloop, Bernhard Schmidt, Marie K. Blicher, Francesca Severino, Grégoire Wuerzner, Alison Taylor, Antoinette Pechère-Bertschi, Fadi Jokhaji, Fadl Elmula M. Fadl Elmula, Jan Rosa, Danuta Czarnecka, Georg Ehret, Thomas Kahan, Jean Renkin, Jirí Widimský, Jr, Lotte Jacobs, Wilko Spiering, Michel Burnier, Patrick B. Mark, Jan Menne, Michael H. Olsen, Peter J. Blankestijn, Sverre Kjeldsen, Michiel L. Bots and Jan A. Staessen

on behalf of the European Network Coordinating research on RENal Denervation (ENCOREd) Consortium

Hypertension. 2014;63:1319-1325; originally published online March 24, 2014;

doi: 10.1161/HYPERTENSIONAHA.114.03194

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2014 American Heart Association, Inc. All rights reserved.

Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hyper.ahajournals.org/content/63/6/1319>

Data Supplement (unedited) at:

<http://hyper.ahajournals.org/content/suppl/2014/03/24/HYPERTENSIONAHA.114.03194.DC1.html>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Hypertension* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:

<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Hypertension* is online at:

<http://hyper.ahajournals.org/subscriptions/>

Eligibility for Renal Denervation Experience at 11 European Expert Centers

Alexandre Persu, Yu Jin, Marie Baelen, Eva Vink, Willemien L. Verloop, Bernhard Schmidt, Marie K. Blicher, Francesca Severino, Grégoire Wuerzner, Alison Taylor, Antoinette Pechère-Bertschi, Fadi Jokhaji, Fadl Elmula M. Fadl Elmula, Jan Rosa, Danuta Czarnecka, Georg Ehret, Thomas Kahan, Jean Renkin, Jiří Widimský Jr, Lotte Jacobs, Wilko Spiering, Michel Burnier, Patrick B. Mark, Jan Menne, Michael H. Olsen, Peter J. Blankestijn, Sverre Kjeldsen, Michiel L. Bots, Jan A. Staessen; on behalf of the European Network Coordinating research on RENal Denervation (ENCOREd) Consortium*

Abstract—Based on the SYMPLICITY studies and CE (Conformité Européenne) certification, renal denervation is currently applied as a novel treatment of resistant hypertension in Europe. However, information on the proportion of patients with resistant hypertension qualifying for renal denervation after a thorough work-up and treatment adjustment remains scarce. The aim of this study was to investigate the proportion of patients eligible for renal denervation and the reasons for noneligibility at 11 expert centers participating in the European Network COordinating Research on renal Denervation in treatment-resistant hypertension (ENCOREd). The analysis included 731 patients. Age averaged 61.6 years, office blood pressure at screening was 177/96 mmHg, and the number of blood pressure-lowering drugs taken was 4.1. Specialists referred 75.6% of patients. The proportion of patients eligible for renal denervation according to the SYMPLICITY HTN-2 criteria and each center's criteria was 42.5% (95% confidence interval, 38.0%–47.0%) and 39.7% (36.2%–43.2%), respectively. The main reasons of noneligibility were normalization of blood pressure after treatment adjustment (46.9%), unsuitable renal arterial anatomy (17.0%), and previously undetected secondary causes of hypertension (11.1%). In conclusion, after careful screening and treatment adjustment at hypertension expert centers, only ≈40% of patients referred for renal denervation, mostly by specialists, were eligible for the procedure. The most frequent cause of ineligibility (approximately half of cases) was blood pressure normalization after treatment adjustment by a hypertension specialist. Our findings highlight that hypertension centers with a record in clinical experience and research should remain the gatekeepers before renal denervation is considered. (*Hypertension*. 2014;63:1319-1325.) • [Online Data Supplement](#)

Key Word: Antihypertensive Agents ■ Hypertension Resistant to Conventional Therapy ■ Sympathetic Denervation

Depending on populations studied and applied methods and definitions, the prevalence of treatment-resistant hypertension varies from 3% to 30%.^{1,2} The SYMPLICITY studies³⁻⁵ demonstrated that in this indication catheter-based endovascular sympathetic renal denervation (RDN) by

means of low-frequency energy is feasible. It entails a 25- to 30-mmHg decrease in office systolic blood pressure, 84% of patients achieving a decrease in office systolic blood pressure of ≥10 mmHg with a rate of procedural adverse events <5% assessed 6 months after RDN.⁴ However, as reviewed

Received January 18, 2014; first decision February 9, 2014; revision accepted February 26, 2014.

From the Pole of the Cardiovascular Research, Institut de Recherche Expérimentale et Clinique (A.P., J. Renkin) and Division of Cardiology, Cliniques Universitaires Saint-Luc (A.P., M.B., F.S., J. Renkin), Université Catholique de Louvain, Brussels, Belgium; Studies Coordinating Centre, Research Unit Hypertension and Cardiovascular Epidemiology, KU Leuven Department of Cardiovascular Sciences, University of Leuven, Leuven, Belgium (Y.J., L.J., J.A.S.); Julius Center for Health Sciences and Primary Care, Utrecht, The Netherlands (M.L.B.); Departments of Vascular Medicine (W.S.), Nephrology (E.V., P.J.B.), and Cardiology (W.V.), University Medical Center Utrecht, Utrecht, The Netherlands; Department of Nephrology and Hypertension, Hannover Medical School, Hannover, Germany (B.S., J.M.); Cardiovascular and Metabolic Preventive Clinic, Department of Endocrinology, Center for Individualized Medicine in Arterial Diseases, Odense University Hospital, Odense, Denmark (M.K.B., M.H.O.); Division of Nephrology, Lausanne University Hospital, Lausanne, Switzerland (G.W., M.B.); BHF Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow, United Kingdom (A.T., P.M.); Hypertension Unit (A.P.-B.) and Division of Cardiology (G.E.), Geneva University Hospitals, Geneva, Switzerland; Karolinska Institute, Department of Clinical Sciences, Danderyd Hospital, Division of Cardiovascular Medicine, Stockholm, Sweden (F.J., T.K.); Department of Cardiology, Ullevål University Hospital, University of Oslo, Oslo, Norway (F.E.M.F.E., S.E.K.); Third Department of Internal Medicine, Centre for Hypertension, General University Hospital, First Faculty of Medicine, Charles University, Prague, Czech Republic (J. Rosa, J.W.); and First Department of Cardiology, Interventional Electrophysiology and Hypertension, Jagiellonian University Medical College, Kraków, Poland (D.C.).

*A list of all European Network Coordinating research on RENal Denervation (ENCOREd) consortium participants is given in the Appendix.

This paper was sent to Robert Carey, Consulting editor, for review by expert referees, editorial decision, and final disposition.

The online-only Data Supplement is available with this article at <http://hyper.ahajournals.org/lookup/suppl/doi:10.1161/HYPERTENSIONAHA.114.03194/-/DC1>.

Correspondence to Jan A. Staessen, Studies Coordinating Centre, Research Unit Hypertension and Cardiovascular Epidemiology, KU Leuven Department of Cardiovascular Sciences, University of Leuven, Kapucijnenvoer 35, Box 7001, BE-3000 Leuven, Belgium. E-mail jan.staessen@med.kuleuven.be

© 2014 American Heart Association, Inc.

Hypertension is available at <http://hyper.ahajournals.org>

DOI: 10.1161/HYPERTENSIONAHA.114.03194

elsewhere,^{6,7} blood pressure decrease observed after RDN is highly variable⁸ and probably much less than previously thought, especially when ambulatory blood pressure monitoring is used.^{8–10} Furthermore, stenosis^{11,12} and damage¹³ of the renal arteries can occur after RDN, and the long-term effect of RDN on renal function is still unclear.^{5,14,15} Hence, until more and better evidence is available, RDN should remain the last resort in patients with truly resistant hypertension.^{6,7} Accordingly, several national and international consensus papers^{16,17} have proposed guidelines for evaluation and management of patients with resistant hypertension before considering RDN. The proportion of patients with truly resistant hypertension eligible for RDN and the reasons of noneligibility after thorough screening and optimization of drug treatment in expert centers remain elusive. In this study, we reviewed the reasons for noneligibility at 11 hypertension expert centers performing RDN for treatment-resistant hypertension and collaborating within the European Network COordinating research on Renal Denervation (ENCOREd).⁸

Methods

Patients

We performed systematic reviews of the literature published elsewhere^{6,7} and identified ENCOREd centers engaging in RDN. At the fourth ENCOREd network meeting, held in Leuven on April 26, 2013, all 12 centers involved in ENCOREd were invited and accepted to take part in this study. Eventually, 1 center was removed from the final analysis because not all information required for the present analysis could be retrieved from the local database. The 11 other contributing centers provided information on 731 consecutive patients referred for RDN. All patients underwent a center-specific 3- to 6-month diagnostic and therapeutic work-up. The eligibility criteria for RDN at the participating centers were similar to those of the SYMPLICITY studies with the following differences: (1) office systolic blood pressure should be ≥ 140 mmHg (3 centers) or ≥ 160 mmHg in patients without diabetes mellitus or ≥ 150 mmHg in patients with diabetes mellitus (3 centers), or the 24-hour systolic blood pressure should be ≥ 130 mmHg, irrespective of the level of office blood pressure (8 centers); (2) patients with white-coat-resistant hypertension were not eligible (8 centers); (3) evaluation of renal artery anatomy by computed tomography or MRI was mandatory (all centers); (4) patients with >2 renal arteries were eligible provided that the length and diameter of the arteries fulfilled the SYMPLICITY criteria (all centers); (5) the lower limit of the estimated glomerular filtration rate (eGFR) using the modification of diet in renal disease formula was 30 mL/min per 1.73 m² (5 centers) or 45 mL/min per 1.73 m² (3 centers); in 1 center, patients were excluded if serum creatinine was >200 μ mol/L and 2 centers did not set limits to eGFR; and (6) drug adherence was assessed according to electronic compliance monitoring (1 center), plasma dosage of drugs and their metabolites (1 center), pharmacy refill (1 center), witnessed drug intake (3 centers), or the Morisky adherence questionnaire (1 center).

Investigators completed a template data sheet for all these patients. It collected information on anthropometric characteristics of the referred patients, their office and ambulatory blood pressure, drug treatment at the first visit, eligibility for RDN according to the SYMPLICITY HTN-2 inclusion and exclusion criteria (E1)⁴ and the centers' own assessment (E2), and reasons of noneligibility. The comprehensive list of noneligibility criteria included older age/frailty; morbid obesity; altered renal function; previous cholesterol emboli or diffuse atherosclerosis; white-coat-resistant hypertension (conventional blood pressure $\geq 140/90$ mmHg but daytime ambulatory blood pressure $<135/85$ mmHg); renal artery stenosis; renal artery stenting; other anatomic reasons precluding RDN; primary aldosteronism; pheochromocytoma; other causes of secondary hypertension; blood pressure controlled or substantially improved after treatment

adjustment, including or not low-dose (25–50 mg) spironolactone; poor drug adherence; and patient or referring physician's refusal. Furthermore, participants had the possibility to add other reasons as free text. For each patient considered noneligible by the center (E2), >1 reason of noneligibility could be provided.

Statistical Methods

We used SAS software (SAS Institute, Cary, NC), version 9.3, for database management and statistical analysis. Continuous data are presented as mean \pm SD and categorical data as frequencies and percentages. We compared means and proportions by a large sample z test and the χ^2 statistic, respectively. We applied a κ statistic to assess agreement between diagnostic criteria. We applied stepwise logistic regression to search for variables associated with an increased likelihood of being eligible for RDN. P values for independent variables to enter and stay the model were set at 0.15. The variables considered for entering into the models were sex, age, body mass index, baseline systolic blood pressure, eGFR, total number of drugs taken, a history of diabetes mellitus, coronary artery disease, stroke, and peripheral artery disease. Odds ratios express the independent increase in risks associated with a 1-SD increase in continuous covariables or a condition of interest (eg, sex). Significance was a 2-sided P value of ≤ 0.05 .

Results

Baseline Characteristics

Information was obtained on 731 subjects. Specialists referred 75.6% of patients (mostly cardiologists and nephrologists), and general practitioners referred 24.4% of patients. Baseline blood pressure, the frequency of comorbidities, drug treatment score, and drug classes (Table 1) were similar to those observed in SYMPLICITY HTN-2.⁴ Notably, at the first visit to the referral center, 14.9% of patients had no drug inhibiting the renin-angiotensin system, 27.2% had no calcium antagonist, and 18.0% had no diuretic (Table 2; Figure S1 in the online-only Data Supplement). Furthermore, 21.5% were prescribed 2 or even 3 drugs inhibiting the renin-angiotensin system. The proportion of patients under spironolactone was low (26.0%) even in patients with eGFR ≥ 45 (28.1%) or ≥ 60 mL/min per 1.73 m² (27.0%). The proportion of patients who had taken spironolactone in the past was 20.9% (95% confidence interval [CI], 17.1%–24.7%), 22.4% (CI, 18.2%–26.6%), and 22.8% (CI, 18.1%–27.5%) in the whole group and in patients with eGFR ≥ 45 or ≥ 60 mL/min per 1.73 m², respectively.

Eligibility Status

Fewer than half of patients were eligible for RDN, both according to the SYMPLICITY HTN-2 criteria (E1, 42.5%; CI, 38.0%–47.0%) and the centers' own judgment (E2, 39.7%; CI, 36.2%–43.2%), with substantial variability between centers (Table 3). In patients with apparently resistant hypertension (office systolic blood pressure ≥ 140 mmHg, despite prescription of ≥ 3 antihypertensive drug classes; $n=601$) and in patients with office systolic blood pressure ≥ 160 mmHg and eGFR ≥ 45 mL/min per 1.73 m² ($n=433$), the proportion of eligible patients was slightly higher (E1, 45.4% and E2, 40.1%; and E1, 60.2% and E2, 48.3%, respectively). Eligibility status was moderately consistent between E1 and E2 (κ coefficient of agreement, 0.49). As may be expected, the proportion of eligible patients was higher in the subset referred by specialists than in patients referred by general practitioners (E1, 47.0% versus 31.6%; $P<0.001$; and E2, 43.6% versus 27.5%; $P<0.001$).

Table 1. Main Characteristics of Patients Referred for RDN According to Eligibility Status

Characteristics	All Patients n=731	E2 Patients n=290 (39.7%)	Non-E2 Patients n=441 (60.3%)
Women, n (%)	336 (46.0)	122 (42.1)	214 (48.5)
Age, y	61.6±12.0	60.0±11.3	62.7±14.4*
White, n (%)	701 (95.9)	276 (95.2)	425 (96.4)
Office systolic pressure, mm Hg	176.7±27.2	183.8±27.6	172.1±25.9†
Office diastolic pressure, mm Hg	96.2±16.8	100.9±17.1	93.2±16.0†
24-h systolic pressure, mm Hg	154.7±20.1	160.7±17.9	149.8±20.5†
24-h diastolic pressure, mm Hg	89.5±14.6	94.4±13.6	85.5±14.2†
No. of drug classes	4.1±1.6	4.3±1.8	3.9±1.5*
Body mass index, kg/m ²	30.0±5.6	29.0±5.8	30.2±5.5
Diabetes mellitus, n (%)	184/668 (27.5)	75/278 (27.0)	109/390 (28.0)
Coronary artery disease, n (%)	134/667 (20.1)	54/277 (19.5)	80/390 (20.5)
Stroke, n (%)	44/457 (9.6)	14/171 (8.2)	30/286 (10.5)
Peripheral artery disease, n (%)	83/665 (12.5)	33/278 (11.9)	50/387 (12.9)
eGFR classes, mL/min per 1.73 m ²	73.8±22.5	76.1±21.7	72.3±22.9‡
	n=686	n=275	n=411
<30	15 (2.2)	2 (0.7)	13 (3.2)
30–45	64 (9.3)	19 (6.9)	45 (11.0)
≥45	607 (88.5)	254 (92.4)	353 (85.9)

Data are mean±SD or number of patients (%). E2 indicates patients eligible for renal denervation according to the centers' own criteria; eGFR, estimated glomerular filtration rate; non-E2, patients ineligible for renal denervation according to the centers' own criteria; and RDN, renal denervation.

Difference between E2 and non-E2 patients: * $P<0.01$, † $P<0.001$, ‡ $P<0.05$. Difference in the eGFR classification between E2 and non-E2: $P=0.017$.

Determinants of Eligibility

Compared with patients considered ineligible by the centers (non-E2), eligible patients (E2) were slightly but significantly younger, had a significantly higher office and ambulatory blood pressure, were taking more drug classes, and had a higher eGFR (Table 1). In a stepwise logistic regression model, age, baseline blood pressure, number of blood pressure-lowering drugs, eGFR, and body mass index were associated with the likelihood of eligibility for RDN. Eligibility for RDN was independently related with 1-SD increase in office systolic blood pressure (odds ratio [OR], 1.87; CI, 1.54–2.25; $P<0.001$), increase in the number of blood pressure-lowering drugs (OR, 1.31; CI, 1.09–1.57; $P=0.004$), and increase in eGFR (OR, 1.25; CI, 1.04–1.51; $P=0.02$), lower age (OR, 0.72; CI, 0.59–0.87; $P=0.001$), and lower body mass index (OR, 0.84; CI, 0.70–1.01; $P=0.064$).

Reasons for noneligibility with a frequency of ≥5% (Figures 1 and 2; Table S1) were (1) controlled or substantially improved blood pressure after treatment adjustment (46.9%), in 49.7% of cases using a regimen including low-dose spironolactone (25–50 mg/d); (2) anatomic reasons (17.0%, renal artery stenosis, history of renal stenting, or other anatomic reasons); (3) secondary hypertension (11.1%, most often primary aldosteronism but also including autonomic dysfunction and hypercortisolism); (4) decreased renal function (9.1%); (5) older age and frailty (7.9%); (6) white-coat-resistant hypertension (7.7%); and (7) poor drug adherence (7.7%).

A nonexhaustive list of other reasons of noneligibility encompassed morbid obesity (n=17); patient (n=10) or referring physician (n=3) refusing the procedure; diffuse atherosclerosis or

previous cholesterol emboli (n=2); proteinuria (n=2); clear cell carcinoma/renal mass (n=2); subarachnoid hemorrhage (n=1) or unstable angina pectoris (n=2) before RDN; nephrotic proteinuria because of light chain disease (n=1); bilateral fibromuscular dysplasia (n=1); hydronephrosis (n=1); primary hyperparathyroidism (n=1); chronic autoimmune disease (n=1); psychiatric illness (n=1); or alcohol abuse (n=1). Of note, in 26.3% of patients deemed ineligible, ≥2 reasons of noneligibility were present.

Discussion

In this multicenter cohort, only ≈40% of patients referred for RDN, mostly by specialists, were eligible for the procedure, according to the SYMPPLICITY HTN-2 criteria, as well as according to the centers' own judgment. These data confirm at the European level the results obtained by Verloop et al¹⁸ in a Dutch single-center cohort, also included in ENCOREd. The proportion of less complex, standard patients with resistant hypertension in whom RDN seems both feasible and justified is probably still an order of magnitude lower (7.5% in patients with resistant hypertension hospitalized for hypertension in a French third referral center; 2.6% according to a recent US chart review)^{2,19} and might further decrease if all patients would be assessed for drug adherence²⁰ or would have received maximal and optimized drug treatment, including low-dose spironolactone whenever possible.¹⁹

As in the Dutch study,¹⁸ patients found to be ineligible for RDN in our current study were older and had lower office and ambulatory blood pressures and medication scores. Although the proportion of patients eligible for RDN was of the same

Table 2. Drug Treatment of Patients Referred for Renal Denervation at the First Visit Before Treatment Adjustment

Drug Treatment	No. of Patients
Total number	692
Antihypertensive drug classes, n	4.1±1.6
Inhibitors of the renin–angiotensin system	
Total	589 (85.1)
Converting enzyme inhibitors (n=418)	180 (43.1)
Angiotensin receptor blockers (n=418)	223 (53.4)
Aliskiren (n=418)	60 (14.4)
≥2 inhibitors of the renin system (n=418)	90 (21.5)
Calcium channel blockers	504 (72.8)
Diuretics	
Total	568 (82.0)
Thiazide (n=418)	279 (66.8)
Loop diuretic (n=418)	120 (28.7)
Spironolactone	180 (26.0)
β-Blockers	498 (71.9)
α-Blockers	141 (20.4)
Centrally acting drugs	175 (25.3)
Vasodilators	42 (6.1)

The Utrecht center did not have detailed information on file on the subclassification of renin system inhibitors or diuretics.

order of magnitude in the Dutch and our study (33.1% versus 39.7%), the leading causes of ineligibility substantially differed. In particular, whereas in the Dutch cohort,¹⁸ successful treatment adjustment excluded only 12% of the referred patients from RDN, in our European cohort, treatment optimization allowed to control blood pressure in ≈50% of noneligible patients, despite the fact that most patients were referred because blood pressure control was considered impossible to achieve, often after years of unsuccessful attempts.

In half of cases (Figure 1), treatment adjustment involved prescription of low-dose spironolactone (25–50 mg/d).

Additional changes might have included prescription of other drug classes, optimization of the dose of diuretics and stepped care combination of diuretics acting at different nephron segments (sequential nephron blockade),²¹ use of more potent or long-acting drugs, and use of double or triple fixed drug combinations to increase drug adherence. One might also hypothesize that consultation in a specialized center and the perspective of use of RDN, an invasive and still exceptional procedure, might have convinced a proportion of patients to adhere more strictly to their drug treatment, which may explain substantial blood pressure improvements observed in some patients despite minimal treatment changes.

Once more, the proportion of patients amenable to blood pressure control might be an underestimation because not all investigators attempted to obtain maximal treatment in all patients. One may expect that the proportion is higher in unselected cohorts of patients with resistant hypertension. These findings are at odds with the frequent belief, held by some investigators and disseminated by the device industry that, for patients with resistant hypertension, per definition, no reasonable further pharmaceutical options are available.²² They highlight the importance of skilful treatment adjustment and the important role of hypertension tertiary referral centers as gatekeepers before RDN is considered.

The effect of treatment optimization should also be interpreted in the light of the drug regimen at the first referral visit, which was neither maximal nor optimal in a substantial proportion of patients. In line with the data published by Hanselin et al,²³ the association of 2 or even 3 renin–angiotensin system blockers, a less efficacious and potentially harmful drug combination,^{24,25} was used in 22% of patients, whereas diuretics (82%) and, notably, calcium antagonists (73%) were still underused. In particular, low-dose spironolactone, a recommended drug in patients with resistant hypertension^{26,27} was prescribed in only 26% of patients (Table 2), even in a subset with normal renal function, which is much higher than in the US database explored by Hanselin et al²³ (5.9%) and slightly better than in 2 other cohorts of patients referred in

Table 3. Proportion of Referred Patients Eligible for RDN After Thorough Work-Up and Treatment Adjustment

Center Name	No. of Patients	Eligible According to SYMPLICITY HTN-2 Criteria (E1) Number (%), 95% Confidence Interval	Eligible According to Centers' Own Criteria (E2) Number (%), 95% Confidence Interval
Lausanne	53	19 (35.9), 23.0–48.8	14 (26.4), 14.5–38.3
Geneva	32	16 (50.0), 32.7–67.3	12 (37.5), 20.7–54.3
Hannover	91	44 (48.4), 38.1–58.7	48 (52.8), 42.5–63.1
Prague	18	14 (77.8), 58.6–97.0	4 (22.2), 3.0–41.4
Oslo	18	10 (55.6), 32.6–78.6	6 (33.3), 11.5–55.1
Stockholm	21	15 (71.4), 52.1–90.7	17 (81.0), 64.2–97.8
Odense	64	7 (10.9), 3.3–18.5	7 (10.9), 3.3–18.5
Glasgow	52	33 (64.5), 51.5–77.5	25 (48.1), 34.5–61.7
Brussels	91	31 (34.1), 24.2–43.8	33 (36.3), 26.4–46.2
Utrecht	274	...	119 (43.4), 37.5–49.3
Krakow	17	5 (29.4), 7.7–51.1	5 (29.4), 7.7–51.1
All centers	731	194/457 (42.5), 38.0–47.1	290/731 (39.7), 36.2–43.2

RDN indicates renal denervation.




Figure 1. Reasons of noneligibility for renal denervation after thorough work-up and treatment optimization in 11 expert hypertension centers. For each patient, several reasons of noneligibility could be applicable (see Figure 2). Only reasons with a frequency of $\geq 5\%$ were plotted.

hypertension expert centers (24%)^{18,19} but clearly remains below expectations. Indeed, although spironolactone was tried in the past but stopped at the time of referral in the expert center in $\approx 20\%$ additional patients, likely because of poor efficacy or drug intolerance, half of patients were never exposed to a mineralocorticoid receptor antagonist.

The frequency of other causes of ineligibility for RDN deserves specific comments. Compared with Simplicity HTN-2,⁴ the proportion of patients excluded for anatomic reasons was similar (17% versus 24%), despite the fact that, in our cohort, patients with multiple renal arteries were considered as eligible, provided the length and diameter of the main

renal arteries were ≥ 20 mm and 4 mm, respectively.⁴ Although the low frequency of white-coat-resistant hypertension (8% versus 38% in the Spanish registry on ambulatory blood pressure monitoring)²⁸ likely reflects preselection by referring physicians, the prevalence of secondary hypertension was similar to that reported in other studies (12%),^{18,29} despite the fact that most patients were referred by specialists.

This study has to be interpreted within the context of its limitations, the most important being the heterogeneity in referral and eligibility criteria, screening, selection, and treatment algorithms between the different centers. Second, the exact nature and dosage of drugs used, proportion of single-pill versus fixed-dose combinations, and the precise strategy used by each center for treatment adjustment were not available. Finally, drug adherence was not evaluated according to standardized methods in all centers. However, our study, the largest conducted thus far, has the advantage of involving multiple centers, thus providing a more representative overview of screening for RDN in European hypertension centers. It also generated information on the effect of treatment adjustment in resistant hypertension. Furthermore, the limitations mentioned previously might only have led to an underestimation of the proportion of ineligible patients, and in particular of patients amenable to blood pressure target after drug treatment optimization, and thus are unlikely to affect the key messages of this study. These are (1) only a minority of patients referred for RDN are eligible for the technique; (2) the most frequent cause of ineligibility (46.9%) was blood pressure normalization after treatment adjustment in the expert center; and (3) in more than half of patients successfully controlled by drug treatment optimization, low-dose spironolactone was part of the therapeutic arsenal.

Perspectives

In view of the persistent uncertainties on the safety and efficacy of RDN—further emphasized by the recent announcement of the failure of SYMPPLICITY HTN-3³⁰ to reach its primary efficacy end point (<http://www.tctmd.com/show.aspx?id=123265>)—difficult to treat patients and patients with resistant hypertension should be thoroughly evaluated in an expert hypertension center before RDN is considered (http://www.bcs.com/pages/news_full.asp?NewsID=19792021). More generally, the availability of RDN as last-line treatment for patients with resistant hypertension is a unique opportunity to revisit the diagnostic and therapeutic algorithms of resistant hypertension and to increase awareness about the best diagnostic and therapeutic strategies in this small subset of patients with hypertension. Finally, although current eligibility criteria are mostly based on SYMPPLICITY HTN-2⁴ and medical common sense, future works should focus on the identification of patients most likely to benefit from RDN, with the lowest probability of adverse events or complications. In particular, identification of predictive factors of blood pressure response post-RDN is a priority. In this respect, our first study on the efficacy of RDN within the ENCOReD network suggested a higher probability of blood pressure response in patients with established cardiovascular disease.⁸ Conversely, higher baseline serum creatinine was associated with lower probability of blood pressure improvement after RDN.⁸




Figure 2. Main reasons of noneligibility of patients with resistant hypertension screened for renal denervation. BP indicates blood pressure; and HT, hypertension.

Although preliminary, these results may pave the way toward identification of responder and nonresponder profiles in currently planned or ongoing randomized controlled studies.

Appendix

Brussels—Marie Baelen, Bernhard Gerber, Sandrine Horman, Joëlle Kefer, Jean-Philippe Lengelé, Jean-Benoit le Polain de Waroux, Jean Renkin, Christophe Scavée, Francesca Severino, and Jean-Louis Vanoverschelde; **Geneva**—Georg Ehret and Antoinette Péchère-Bertschi; **Glasgow**—Collin Berry, Adrian Brady, Christian Delles, Anna Dominiczak, Marie Freel, Alan Jardine, Jon Moss, Scot Muir, Patrick Mark, Sandosh Padmanabhan, and Giles Roditi; **Hannover**—Johann Bauersachs, Julia Brinkmann, Hermann Haller, Karsten Heusser, Jens Jordan, Gunnar Klein, Jan Menne, Bernhard Schmidt, and Jens Tank; **Krakow**—D. Czarnecka, Marek Jastrzębski, and Katarzyna Styczkiewicz; **Lausanne**—Michel Burnier and Grégoire Wuerzner; **Leuven**—Kei Asayama, Yumei Gu, Asuza Hashimoto, Lotte Jacobs, Yu Jin, Tatiana Kuznetsova, Yanping Liu, Lutgarde Thijs, and Jan A Staessen; **Odense**—Maria Blicher, Henning Beck-Nielsen, Poul Flemming Højlund-Carlsen, and M. Olsen; **Oslo**—Magne Brekke, Kristian Engeseth, Fadl Elmula M. Fadl Elmula, Eigil Fossum, Eivind Gjønness, Ulla Hjørnholm, Pavel Hoffmann, Aud Høiegggen, Vibeke Kjær, Sverre E. Kjeldsen, Anne C.K. Larstorp, Oliver Meyerdericks, Ingrid Os, Morten Rostrup, and Aud Stenehjem; **Prague**—Jan Rosa, Ondrej Petrak, Tomas Zelinka, Branislav Strauch, Karol Curila, Petr Tousek, Jiří Widimský and Petr Widimský; **Stockholm**—Fadi Jokhaji, Riker Lander, Thomas Kahan, and Jonas Spaak; **Utrecht**—Peter J. Blankestijn, Michiel L. Bots, Pieter A. Doevendans, Maarten B. Rookmaaker, Wilko Spiering, Willemien L. Verloop, Eva E. Vink, Michiel Voskuil, and Evert-Jan Vonken.

Acknowledgments

We gratefully acknowledge the expert assistance of Sandra Covens and Annick de Soete (Studies Coordinating Centre, Leuven, Belgium).

Sources of Funding

The European Union (grants IC15-CT98-0329-EPOGH, LSHM-CT-2006-037093-InGenious HyperCare, HEALTH-2007-2.1.1-2-HyperGenes, HEALTH-2011.2.4.2-2-EU-MASCARA, HEALTH-F7-305507 HOMAGE, and the European Research Council Advanced Researcher Grant-2011-294713-EPORE) and the Ministry of the Flemish Community, Brussels, Belgium (grants G.0734.09, G.0881.13, and G.088013N) supported the Studies Coordinating Center, Leuven, Belgium. Other sources of funding include the Swiss National Foundation (grant number 140331). E. Vink was supported by a grant of the Dutch Kidney Foundation (IP10-15).

Disclosures

P.J. Blankestijn received unrestricted research grants from Medtronic Inc and speaker and consultancy fees from Medtronic Inc and St. Jude Medical Inc. The other authors report no conflicts.

References

1. Staessen JA, Kuznetsova T, Stolarz K. Hypertension prevalence and stroke mortality across populations. *JAMA*. 2003;289:2420–2422.
2. Hayek SS, Abdou MH, Demoss BD, Legaspi JM, Veledar E, Deka A, Krishnan SK, Wilmut KA, Patel AD, Kumar VR, Devireddy CM. Prevalence of resistant hypertension and eligibility for catheter-based renal denervation in hypertensive outpatients. *Am J Hypertens*. 2013;26:1452–1458.
3. Krum H, Schlaich M, Whitbourn R, Sobotka PA, Sadowski J, Bartus K, Kapelak B, Walton A, Sievert H, Thambar S, Abraham WT, Esler M. Catheter-based renal sympathetic denervation for resistant hypertension: a multicentre safety and proof-of-principle cohort study. *Lancet*. 2009;373:1275–1281.
4. Symplicity HTN-2 Investigators. Renal sympathetic denervation in patients with treatment-resistant hypertension (The Symplicity HTN-2 Trial): a randomised controlled trial. *Lancet*. 2010;376:1903–1909.
5. Symplicity HTN-1 Investigators. Catheter-based renal sympathetic denervation for resistant hypertension. Durability of blood pressure reduction out to 24 months. *Hypertension*. 2011;57:911–917.
6. Persu A, Renkin J, Thijs L, Staessen JA. Renal denervation: ultima ratio or standard in treatment-resistant hypertension. *Hypertension*. 2012;60:596–606.
7. Persu A, Renkin J, Asayama K, O'Brien E, Staessen JA. Renal denervation in treatment-resistant hypertension: the need for restraint and more and better evidence. *Expert Rev Cardiovasc Ther*. 2013;11:739–749.
8. Persu A, Jin Y, Azizi M, et al. Blood pressure changes after renal denervation at 10 European expert centers. *J Hum Hypertens*. 2014;28:150–156.
9. Mahfoud F, Ukena C, Schmieder RE, et al. Ambulatory blood pressure changes after renal sympathetic denervation in patients with resistant hypertension. *Circulation*. 2013;128:132–140.
10. Howard JP, Nowbar AN, Francis DP. Size of blood pressure reduction from renal denervation: insights from meta-analysis of antihypertensive drug trials of 4,121 patients with focus on trial design: the CONVERGE report. *Heart*. 2013;99:1579–1587.
11. Vonend O, Antoch G, Rump LC, Blondin D. Secondary rise in blood pressure after renal denervation. *Lancet*. 2012;380:778.
12. Kaltenbach B, Id D, Franke JC, Sievert H, Hennersdorf M, Maier J, Bertog SC. Renal artery stenosis after renal sympathetic denervation. *J Am Coll Cardiol*. 2012;60:2694–2695.
13. Templin C, Jaguszewski M, Ghadri JR, Sudano I, Gachwiler R, Hellermann JP, Schoenenberger-Berzins R, Landmesser U, Erne P, Noll G, Lüscher TF. Vascular lesions induced by renal nerve ablation as assessed by optical coherence tomography: pre- and post-procedural comparison with the Simplicity catheter system and the EnligHTN multi-electrode renal denervation catheter. *Eur Heart J*. 2013;34:2141–2148, 2148b.
14. Petidis K, Anyfanti P, Doumas M. Renal sympathetic denervation: renal function concerns. *Hypertension*. 2011;58:1–19.
15. Worthley SG, Tsioufis CP, Worthley MI, Sinhal A, Chew DP, Meredith IT, Malaipayan Y, Papademetriou V. Safety and efficacy of a multi-electrode renal sympathetic denervation system in resistant hypertension: the EnligHTN I trial. *Eur Heart J*. 2013;34:2132–2140.
16. Schmieder RE, Redon J, Grassi G, Kjeldsen SE, Mancia G, Narkiewicz K, Parati G, Ruilope L, van de Borne P, Tsioufis C. ESH position paper: renal denervation - an interventional therapy of resistant hypertension. *J Hypertens*. 2012;30:837–841.
17. Mahfoud F, Lüscher TF, Andersson B, et al; European Society of Cardiology. Expert consensus document from the European Society of Cardiology on catheter-based renal denervation. *Eur Heart J*. 2013;34:2149–2157.
18. Verloop WL, Vink EE, Voskuil M, Vonken EJ, Rookmaaker MB, Bots ML, Doevendans PA, Blankestijn PJ, Spiering W. Eligibility for percutaneous renal denervation: the importance of a systematic screening. *J Hypertens*. 2013;31:1662–1668.
19. Savard S, Frank M, Bobrie G, Plouin PF, Sapoval M, Azizi M. Eligibility for renal denervation in patients with resistant hypertension: when enthusiasm meets reality in real-life patients. *J Am Coll Cardiol*. 2012;60:2422–2424.
20. Fadl Elmula FE, Hoffmann P, Fossum E, Brekke M, Gjønness E, Hjørnholm U, Kjær VN, Rostrup M, Kjeldsen SE, Os I, Stenehjem AE, Høiegggen A. Renal sympathetic denervation in patients with treatment-resistant hypertension after witnessed intake of medication before qualifying ambulatory blood pressure. *Hypertension*. 2013;62:526–532.
21. Bobrie G, Frank M, Azizi M, Peyrard S, Boutouyrie P, Chatellier G, Laurent S, Menard J, Plouin PF. Sequential nephron blockade versus sequential renin-angiotensin system blockade in resistant hypertension: a prospective, randomized, open blinded endpoint study. *J Hypertens*. 2012;30:1656–1664.
22. Brandt MC, Reda S, Mahfoud F, Lenski M, Böhm M, Hoppe UC. Effects of renal sympathetic denervation on arterial stiffness and central hemodynamics in patients with resistant hypertension. *J Am Coll Cardiol*. 2012;60:1956–1965.
23. Hanselin MR, Saseen JJ, Allen RR, Marrs JC, Nair KV. Description of antihypertensive use in patients with resistant hypertension prescribed four or more agents. *Hypertension*. 2011;58:1008–1013.
24. Mann JF, Schmieder RE, McQueen M, et al; ONTARGET Investigators. Renal outcomes with telmisartan, ramipril, or both, in people at high vascular risk (the ONTARGET study): a multicentre, randomised, double-blind, controlled trial. *Lancet*. 2008;372:547–553.

25. Parving HH, Brenner BM, McMurray JJ, de Zeeuw D, Haffner SM, Solomon SD, Chaturvedi N, Persson F, Desai AS, Nicolaidis M, Richard A, Xiang Z, Brunel P, Pfeffer MA; ALTITUDE Investigators. Cardiorenal end points in a trial of aliskiren for type 2 diabetes. *N Engl J Med*. 2012;367:2204–2213.
26. Václavík J, Sedlák R, Plachy M, Navrátil K, Plásek J, Jarkovsky J, Václavík T, Husár R, Kociánová E, Táborský M. Addition of spironolactone in patients with resistant arterial hypertension (ASPIRANT): a randomized, double-blind, placebo-controlled trial. *Hypertension*. 2011;57:1069–1075.
27. Oxlund CS, Henriksen JE, Tarnow L, Schousboe K, Gram J, Jacobsen IA. Low dose spironolactone reduces blood pressure in patients with resistant hypertension and type 2 diabetes mellitus: a double blind randomized clinical trial. *J Hypertens*. 2013;31:2094–2102.
28. de la Sierra A, Segura J, Banegas JR, Gorostidi M, de la Cruz JJ, Oliveras A, Ruilope LM. Clinical features of 8295 patients with resistant hypertension classified on the basis of ambulatory blood pressure monitoring. *Hypertension*. 2011;57:889–890.
29. Pedrosa RP, Drager LF, Gonzaga CC, Sousa MG, de Paula LKG, Amaro ACS, Amodeo C, Bortolotto LA, Krieger EM, Bradley TD, Lorenzi-Filho G. The most common secondary cause of hypertension associated with resistant hypertension. *Hypertension*. 2011;58:811–817.
30. Kandzari DE, Bhatt DL, Sobotka PA, O'Neill WW, Esler M, Flack JM, Katzen BT, Leon MB, Massaro JM, Negoita M, Oparil S, Rocha-Singh K, Straley C, Townsend RR, Bakris G. Catheter-based renal denervation for resistant hypertension: rationale and design of the SYMPPLICITY HTN-3 Trial. *Clin Cardiol*. 2012;35:528–535.

Novelty and Significance

What Is New?

- In a multicenter European cohort of 731 patients referred for renal denervation to leading hypertension centers, only ~40% qualified.
- The most frequent cause of ineligibility (46.9%) was blood pressure normalization after treatment adjustment in the expert center, removing the indication for renal denervation.
- Other reasons of ineligibility included incompatible renal artery anatomy (17.0%) and secondary hypertension (11.1%).

What Is Relevant?

- The diagnosis of resistant hypertension requires out-of-the-office blood pressure monitoring.
- Management of resistant hypertension by combining antihypertensive drugs in optimal doses and combinations is substandard in Europe, highlighting the need for continuing education of physicians.
- Renal denervation remains a last-resort approach that should only be

applied at expert hypertension centers in a context of clinical research. The failure of SYMPPLICITY HTN-3 to reach its primary end point of efficacy supports this contention.

- Hypertension expert centers should be the gatekeepers to give patients access to renal denervation.

Summary

This study showed that in 731 patients representative for clinical practice at 11 European hypertension expert centers (mean age, 61.6 years; office blood pressure, 177/96 mm Hg, number of drugs, 4.1), only 39.7% of patients referred for renal denervation qualified. The main reasons of noneligibility were normalization of blood pressure after treatment adjustment (46.9%), incompatible renal artery anatomy (17.0%), and previously undetected secondary hypertension (11.1%).

HYPERTENSION

Eligibility for Renal Denervation: Experience at 11 European Expert Centers

Eligibility for Renal Denervation

ONLINE SUPPLEMENT

Alexandre Persu, Yu Jin, Marie Baelen, Eva Vink, Willemien Verloop, Bernhard Schmidt, Marie K. Blicher, Francesca Severino, Grégoire Wuerzner, Alison Taylor, Antoinette Pechère-Bertschi, Fadi Jokhaji, Fadi Elmula M. Fadi Elmula, Jan Rosa, Danuta Czarnecka, Georg Ehret, Thomas Kahan, Jean Renkin, Jiří Widimský Jr., Lotte Jacobs, Wilko Spiering, Michel Burnier, Patrick B. Mark, Jan Menne, Michael H. Olsen, Peter J. Blankestijn, Sverre Kjeldsen, Michiel L. Bots, Jan A. Staessen, on behalf of the European Network Coordinating research on REnal Denervation (ENCOREd) consortium

Word Counts: Manuscript 5595; Abstract: 247;

Number: 30 References; 3 Tables; 2 Figures

Correspondence:

Jan A. Staessen, MD, PhD,
Research Unit Hypertension and Cardiovascular Epidemiology,
KU Leuven Department of Cardiovascular Sciences,
University of Leuven,
Kapucijnenvoer 35, Box 7001,
BE-3000 Leuven, Belgium

Telephone: +31-43-3388-2374 (office)
Facsimile: +32-43-388-4128 (office)
email: jan.staessen@med.kuleuven.be

Table S1 Reasons of non-eligibility for renal denervation after thorough work-up and treatment adaptation in the centers (E2) (starts)

Center name	Non-E2 patients	Blood pressure controlled after treatment adjustment			Anatomical reasons			
		Spirolactone given	Spirolactone not given	Total	Renal artery stenosis	Renal stenting	Other	Total
Lausanne	39	8 (20.5)	8 (20.5)	16 (41.0)	2 (5.1)	1 (2.6)	4 (10.3)	5 (12.8)
Geneva	20	4 (20.0)	4 (20.0)	8 (40.0)	0	0	0	0
Hannover	43	22 (51.2)	17 (39.5)	39 (90.7)	1 (2.3)	0	0	1 (2.3)
Prague	14	1 (7.1)	7 (50.0)	8 (57.1)	0	0	0	0
Oslo	12	0	1 (8.3)	1 (8.3)	0	0	1 (8.3)	1 (8.3)
Stockholm	4	0	2 (50.0)	2 (50.0)	0	1 (25.0)	0	1 (25.0)
Odense	57	17 (29.8)	22 (38.6)	39 (68.4)	4 (7.0)	0	3 (5.3)	7 (12.3)
Glasgow	27	3 (11.1)	2 (7.4)	5 (18.5)	2 (7.7)	0	10 (37.0)	12 (44.4)
Brussels	58	20 (34.5)	12 (20.7)	32 (55.2)	4 (6.9)	5 (8.6)	10 (17.2)	17 (29.3)
Utrecht	155	NA	NA	54 (34.8)	14 (9.0)	4 (2.6)	5 (3.2)	23 (14.8)
Krakow	12	1 (8.3)	2 (16.7)	3 (25.0)	3 (25.0)	0	5 (41.7)	8 (66.7)
11 centers	441	76	77	207 (46.9)	30 (6.8)	11 (2.5)	38 (8.6)	75 (17.0)

NA indicates information not available. Data are number of patients (%). The table includes reasons for non-eligibility with a frequency of at least 5%. Numbers do not add up, because patients could have more than one reason making them ineligible.

Table S1 Reasons of non-eligibility for renal denervation after thorough work-up and treatment adaptation in the centers (E2) (continued)

Center name	Secondary hypertension			Reduced renal function	Older age/frailty	White-coat hypertension	Poor drug adherence
	Primary aldosteronism	Other cause	Total				
Lausanne	7 (18.0)	1 (2.6)	8 (20.5)	9 (23.1)	12 (30.8)	1 (2.6)	3 (12.5)
Geneva	0	0	0	2 (10.0)	1 (5.0)	6 (30.0)	3 (15.0)
Hannover	0	0	0	0	2 (4.7)	0	3 (7.0)
Prague	4 (28.6)	0	4 (28.6)	0	0	7 (50.0)	3 (21.4)
Oslo	0	0	0	0	0	5 (41.7)	0
Stockholm	0	1 (25.0)	1 (25.0)	1 (25.0)	0	0	1 (25.0)
Odense	1 (1.8)	0	1 (1.8)	13 (22.8)	9 (15.9)	6 (10.5)	4 (7.0)
Glasgow	0	3 (11.1)	3 (11.1)	2 (7.4)	3 (11.1)	2 (7.4)	2 (7.4)
Brussels	0	0	0	12 (20.7)	8 (13.8)	6 (10.3)	0
Utrecht	25 (16.1)	4 (2.6)	29 (18.7)	NA	NA	NA	9 (6%)
Krakow	0	3 (25.0)	3 (25.0)	1 (8.3)	0	1 (8.3)	6 (50%)
11 centers	37 (8.4)	12 (2.7)	49 (11.1)	40 (9.1)	35 (7.9)	34 (7.7)	34 (7.7)

Data are number of patients (%). Reasons of non-eligibility declared in < 5% of cases are not indicated. NA: not available.




Figure S1 Proportion of prescribed antihypertensive drug classes in patients referred for renal denervation, before treatment adjustment in the 11 expert centers. RAS inhibitors refer to renin angiotensin inhibitors (includes angiotensin converting enzyme inhibitors, angiotensin II receptor antagonists and the direct renin inhibitor aliskiren). CCBs refers to calcium channel blockers.